

Processes of r^{th} largest

Boris Buchmann¹ · Ross Maller¹ ·
Sidney I. Resnick² 

Received: 11 May 2017 / Revised: 4 December 2017 / Accepted: 31 January 2018
© Springer Science+Business Media, LLC, part of Springer Nature 2018

Abstract For integers $n \geq r$, we treat the r th largest of a sample of size n as an \mathbb{R}^∞ -valued stochastic process in r which we denote as $\mathbf{M}^{(r)}$. We show that the sequence regarded in this way satisfies the Markov property. We go on to study the asymptotic behavior of $\mathbf{M}^{(r)}$ as $r \rightarrow \infty$, and, borrowing from classical extreme value theory, show that left-tail domain of attraction conditions on the underlying distribution of the sample guarantee weak limits for both the range of $\mathbf{M}^{(r)}$ and $\mathbf{M}^{(r)}$ itself, after norming and centering. In continuous time, an analogous process $\mathbf{Y}^{(r)}$ based on a two-dimensional Poisson process on $\mathbb{R}_+ \times \mathbb{R}$ is treated similarly, but we note that the continuous time problems have a distinctive additional feature: there are always infinitely many points below the r th highest point up to time t for any $t > 0$. This necessitates a different approach to the asymptotics in this case.

This research was initiated and partially supported by ARC grants DP1092502 and DP160104737. S. Resnick also received significant support from US Army MURI grant W911NF-12-1-0385 to Cornell University; Resnick gratefully acknowledges hospitality, administrative support and space during several visits to the Research School of Finance, Actuarial Studies & Statistics, Australian National University.

✉ Sidney I. Resnick
sir1@cornell.edu

Boris Buchmann
Boris.Buchmann@anu.edu.au

Ross Maller
Ross.Maller@anu.edu.au

¹ Research School of Finance, Actuarial Studies & Statistics, Australian National University, Canberra, ACT 0200, Australia

² School of Operations Research and Information Engineering, Cornell University, Ithaca, NY 14853, USA

Keywords Extremes · Domain of attraction · Markov property · Extremal process

AMS 2000 Subject Classifications Primary—60F05, 60G70, 60K99;
Secondary—60G55, 60G51

1 Introduction

In this paper we consider Markovian and other properties of the order statistics of independent identically distributed (iid) random variables (rvs) in discrete time, and of extremal processes in continuous time. Although venerable these are important issues and research continues to throw up significant new aspects. As a starting point let $M_n^{(r)}$ be the r th largest among iid random variables X_1, \dots, X_n with cdf F . (Precise specifications of the order statistics will be given later.) It is known (Arnold et al. 1984) that for fixed n , the *finite* sequence $(M_n^{(r)})_{r=1,2,\dots,n}$ is Markov if and only if F is continuous on (ℓ_F, u_F) , where ℓ_F and u_F are the left and right extremes of F . We investigate the infinitely many order statistics $(M_n^{(r)}, n \geq r)$ for sample sizes beyond r , and further, derive properties of the whole collection $\{\mathbf{M}^{(r)} = (M_n^{(r)}, n \geq r); r \geq 1\}$, considered as an \mathbb{R}^∞ -valued stochastic process. Apart from their intrinsic interest, our results relate a number of areas and techniques.

We begin in Section 2 by setting up the notation required for, then proving, the Markov property, that the conditional distribution of the infinite sequence

$$\mathbf{M}^{(r+1)} = (M_{r+1}^{(r+1)}, M_{r+2}^{(r+1)}, \dots),$$

knowing all values

$$\mathbf{M}^{(1)} = (M_1^{(1)}, M_2^{(1)}, \dots), \mathbf{M}^{(2)} = (M_2^{(2)}, M_3^{(2)}, \dots), \dots, \mathbf{M}^{(r)} = (M_r^{(r)}, M_{r+1}^{(r)}, \dots),$$

is the same as the conditional distribution knowing only $\mathbf{M}^{(r)}$. No continuity assumptions on F are required for this.

In Section 3 we turn to an investigation of asymptotic properties of the collection $\mathbf{M}^{(r)}$, for large values of r . The weak convergence of $\mathbf{M}^{(r)}$, after norming and centering, is related to domain of attraction theory for the *minimum* of an iid sequence of rvs. A key tool in these proofs is Ignatov's (Ignatov 1986) theorem showing that the r -records of an iid sequence are points of a Poisson random measure.

This study is continued in Section 4 for continuous time r th-order extremal processes. Some notable differences between the discrete and continuous time situations emerge here. In particular, unlike in the discrete case, in the continuous time case there are always infinitely many points below the currently considered order statistic, and thus the convergence criterion has to be modified. Section 5 concludes the paper with some modest final thoughts and open problems.

We conclude the present section by mentioning previous and related work. For alternative proofs and other background on Ignatov's (1977) theorem see Ignatov (1986), Stam (1985), Goldie and Rogers (1984), Engelen et al. (1988), and Resnick (2008). Other treatments of the Markov structure of the finite sequence $(M_n^{(r)})_{r=1,2,\dots,n}$ are in (Goldie and Maller 1999; Rüschendorf 1985; Cramer and Tran

2009) and (Rüschendorf 1985; Cramer and Tran 2009) show that $(M_n^{(r)})_{r=1,2,\dots,n}$ is Markov if information on tied values is incorporated into the sequence. For background on continuous time extremal processes we refer to (Resnick 1974, 1975, 2008; Resnick and Rubinovitch 1973). Additional references are given throughout the text.

2 Markov property of higher order extremal processes with discrete indexing

2.1 Notation and indexing

The statement and proof of the Markov property requires precise and detailed notation so that we keep track of infinite sequences indexed by r where the first members are being moved further out as r increases. To cope with this we use the idea of shifted sequences, with first members replaced by $-\infty$.

To see how this works, set $\mathbb{N} = \{1, 2, 3, \dots\}$, $\mathbb{R}_{-\infty} := \mathbb{R} \cup \{-\infty\} = [-\infty, \infty)$, and conventions $\sum \emptyset = 0$, $\prod \emptyset = 1$, $\pm\infty \times 0 = 0$. Sequence space is $\mathbb{R}_{-\infty}^{\mathbb{N}} := \{\mathbf{x} = (x_n) : x_n \in \mathbb{R}_{-\infty}, n \in \mathbb{N}\}$ endowed with the Borel field associated with the product topology and $\mathbb{R}_{-\infty}^{\mathbb{N}, \uparrow} = \{\mathbf{x} = (x_n) \in \mathbb{R}_{-\infty}^{\mathbb{N}} : x_n \leq x_{n+1}, n \in \mathbb{N}\}$ is the subset of nondecreasing sequences. The partial maxima operator $\bigvee : \mathbb{R}_{-\infty}^{\mathbb{N}} \rightarrow \mathbb{R}_{-\infty}^{\mathbb{N}, \uparrow}$ maps a given sequence $\mathbf{x} = (x_n)_n \in \mathbb{R}_{-\infty}^{\mathbb{N}}$ to its associated sequence of partial maxima $\bigvee \mathbf{x} := (\vee\{x_1, \dots, x_n\})_n$. For $n \in \mathbb{N}$, $y_n^{(1)} \geq y_n^{(2)} \geq \dots \geq y_n^{(n)}$ denotes the order statistics associated with (possibly extended) real numbers $y_1, \dots, y_n \in \mathbb{R}_{-\infty}$.

For a given sequence $\mathbf{x} \in \mathbb{R}_{-\infty}^{\mathbb{N}}$ and $r \in \mathbb{N}$, $n \geq r$, let $m_n^{(r)}$ be the r th largest of x_1, \dots, x_n , arranged in lexicographical order in case of ties. Then set

$$x_n^{(r)} = \begin{cases} -\infty, & \text{if } n < r; \\ m_n^{(r)}, & \text{if } n \geq r. \end{cases}$$

The *extremal sequence of order r associated with \mathbf{x}* is the sequence $\mathbf{x}^{(r)} \in \mathbb{R}_{-\infty}^{\mathbb{N}, \uparrow}$, with finite elements $x_n^{(r)}$ augmented with $-\infty$ as follows:

$$\mathbf{x}^{(r)} = (\underbrace{-\infty, \dots, -\infty}_{r-1 \text{ entries}}, m_n^{(r)}, n \geq r). \quad (2.1)$$

Write $\mathbf{x}^{(0)} := \mathbf{x}$ for the extremal sequence of zero order. The extremal sequence of unit order equals the partial maximum sequence: $\mathbf{x}^{(1)} = \bigvee \mathbf{x}$.

For a sequence $\mathbf{x} = (x_n)_n \in \mathbb{R}_{-\infty}^{\mathbb{N}}$ the *shifted sequence* $\mathbf{x}_{\mathcal{R}}$ is $\mathbf{x}_{\mathcal{R}} = (-\infty, \mathbf{x}) \in \mathbb{R}_{-\infty}^{\mathbb{N}}$, so that we append $-\infty$ in front of \mathbf{x} . For two sequences $\mathbf{x} = (x_n)_n$, $\mathbf{y} = (y_n)_n \in \mathbb{R}_{-\infty}^{\mathbb{N}}$, let

$$\mathbf{x}_{\mathcal{R}} \wedge \mathbf{y} := \{(-\infty)\mathbf{1}_{n=1} + (x_{n-1} \wedge y_n)\mathbf{1}_{n>1}\}_n \in \mathbb{R}_{-\infty}^{\mathbb{N}}$$

be the componentwise minimum of \mathbf{x} and \mathbf{y} , taken after shifting \mathbf{x} to the right with proper augmentation with $-\infty$. Thus, componentwise, when $\mathbf{x} = (x_1, x_2, \dots)$ and

$\mathbf{y} = (y_1, y_2, \dots)$, we have

$$\mathbf{x}_{\mathcal{R}} = (-\infty, x_1, x_2, \dots) \quad \text{and} \quad \mathbf{x}_{\mathcal{R}} \wedge \mathbf{y} = (-\infty, x_1 \wedge y_2, x_2 \wedge y_3, \dots).$$

In Theorem 2.1, we will show a Markov property for the r th largest of an iid sequence, and since recursions are an effective tool for proving a sequence of random elements is Markovian, we first prove a preliminary result focussing on properties of the shifted sequences.

Proposition 2.1 *For $r \in \mathbb{N}$, we have the identity,*

$$\mathbf{x}^{(r+1)} = (\mathbf{x}^{(r)}_{\mathcal{R}} \wedge \mathbf{x}) \quad (2.2)$$

or in component form,

$$x_n^{(r+1)} = \bigvee_{j=r+1}^n x_{j-1}^{(r)} \wedge x_j, \quad r \in \mathbb{N}, n \geq r+1, \quad (2.3)$$

with both sides taken as $-\infty$ for $1 \leq n \leq r$.

Proof Fix an integer r and we prove (2.3) by induction on n . The base of the induction is $n = r+1$ and the left side of Eq. 2.3 is $x_{r+1}^{(r+1)} = \bigwedge_{i=1}^{r+1} x_i$. The right side is $x_r^{(r)} \wedge x_{r+1} = \bigwedge_{i=1}^{r+1} x_i$. So Eq. 2.3 is proved for $n = r+1$.

As an induction hypothesis, assume (2.3) is true for $n = r+p$ for $p \geq 1$ and we verify (2.3) to be true for $n = r+p+1$. The left side of Eq. 2.3 for $n = r+p+1$ is $x_{r+p+1}^{(r+1)} =: LHS$. The right side is

$$RHS := \bigvee_{j=r+1}^{r+p+1} \left(\bigwedge_{j-1}^{(r)} \wedge x_j \right) = \bigvee_{j=r+1}^{r+p} \left(x_{j-1}^{(r)} \wedge x_j \right) \vee \left(x_{r+p}^{(r)} \wedge x_{r+p+1} \right)$$

and from the induction hypothesis this is equal to

$$x_{r+p}^{(r+1)} \vee \left(x_{r+p}^{(r)} \wedge x_{r+p+1} \right). \quad (2.4)$$

Now consider cases:

Case (a) $x_{r+p+1} > x_{r+p}^{(r)}$ Then $x_{r+p}^{(r)} = x_{r+p+1}^{(r+1)}$, so $RHS = x_{r+p}^{(r+1)} \vee x_{r+p}^{(r)} = x_{r+p}^{(r)} = LHS$.

Case (b) $x_{r+p}^{(r+1)} \leq x_{r+p+1} \leq x_{r+p}^{(r)}$ The term in parentheses on the right side of Eq. 2.4 then is

$$x_{r+p}^{(r)} \wedge x_{r+p+1} = x_{r+p+1} = x_{r+p+1}^{(r+1)}$$

and thus

$$RHS = x_{r+p}^{(r+1)} \vee x_{r+p+1}^{(r+1)} = x_{r+p+1}^{(r+1)} = LHS.$$

Case (c) $x_{r+p+1} < x_{r+p}^{(r+1)}$ In this case we have

$$RHS = x_{r+p}^{(r+1)} \vee (x_{r+p}^{(r)} \wedge x_{r+p+1} = x_{r+p}^{(r+1)} \vee x_{r+p+1} = x_{r+p}^{(r+1)})$$

because $x_{r+p+1} < x_{r+p}^{(r+1)}$. It follows that $x_{r+p+1}^{(r+1)} = LHS$.

The three cases exhaust the possibilities and this completes the induction argument. \square

2.2 The iid setting

Now we add the randomness. Let $X = (X_n)_n \in \mathbb{R}^{\mathbb{N}}$ be an iid sequence of rvs in \mathbb{R} with cdf F and set $X^{(0)} = X$. Then for $r \in \mathbb{N}$ the r -th order extremal process is the augmented sequence $X^{(r)} = (X_n^{(r)})_{n \in \mathbb{N}}$ in $\mathbb{R}_{-\infty}^{\mathbb{N}}$ constructed as in Eq. 2.1; specifically,

$$X^{(r)} = (\underbrace{-\infty, \dots, -\infty}_{r-1 \text{ entries}}, M_n^{(r)}, n \geq r), \quad (2.5)$$

where the $M_n^{(r)}$ are the order statistics of X_1, X_2, \dots, X_n defined lexicographically as for the $m_n^{(r)}$ in Eq. 2.1. Note that $X^{(1)} = \bigvee X^{(0)} = \bigvee X$ is the sequence of partial maxima associated with X .

To think about the Markov property for $(X^{(r)}, r \geq 1)$, we imagine conditioning on the monotone sequence $X^{(r)} = \mathbf{x}^{(r)}$. For indices where the sequence $\mathbf{x}^{(r)}$ is a constant, say x , the structure of $X^{(r+1)}$ should be as if we construct the maximum sequence from repeated observations from the conditional distribution of $(X_1 | X_1 \leq x)$. See Fig. 1. The following construction makes this precise.

Let $U = (U_{r,n})_{n,r \in \mathbb{N}}$ be an iid array of uniform r.v.'s in $(0, 1)$. Assume $X = X^{(0)}$ and U are independent random elements. For $m \in \mathbb{R}$ with $F(m) > 0$ the left-continuous inverse $u \rightarrow F^{\leftarrow}(u|m)$ of the conditional cdf $x \rightarrow F(x|m) := P(X_1 \leq$




Fig. 1 Blue dotted lines track height of current maximum process $\mathbf{M}^{(1)}$ generated by vertical lines. The red dotted line tracks the second maximum process $\mathbf{M}^{(2)}$. Note that a jump in $\mathbf{M}^{(1)}$ can affect $\mathbf{M}^{(2)}$ as seen at $n = 6$ where X_4 becomes the new value of $\mathbf{M}^{(2)}$. During intervals where the blue dotted line is constant, $\mathbf{M}^{(2)}$ is obtained by sampling from the distribution conditional on the sampled value being less than the blue height. The range of $\mathbf{M}^{(1)}$ consists of blue tick-marks on y-axis and the range of $\mathbf{M}^{(2)}$ is the red ticks

$x|X_1 \leq m$) is well-defined; otherwise, if $F(m) = 0$ set $F^{\leftarrow}(u|m) = \mathbf{1}_{m>0}$ with $F^{\leftarrow}(u|-\infty) \equiv 0$.

For $r \in \mathbb{N} = \{1, 2, \dots\}$ introduce two sequences $\widehat{X}_{\uparrow+1} = (\widehat{X}_{\uparrow+1,n})_n$ and $\widetilde{X}_{\uparrow+1} = (\widetilde{X}_{\uparrow+1,n})_n$. For the first, we have for $n = 1$ that $\widehat{X}_{\uparrow+1,1} := X_1^{(1)} = X_1$ and, for $n \geq 2$,

$$\widehat{X}_{\uparrow+1,n} := \begin{cases} F^{\leftarrow}(U_{r,n}|X_n^{(r)}) \prod_{1 \leq k \leq r} \mathbf{1}_{X_n^{(k)} = X_{n-1}^{(k)}}, & \text{if } X_n^{(r)} = X_{n-1}^{(r)}, \\ \sum_{k=1}^r X_n^{(k)} \mathbf{1}_{X_n^{(k)} > X_{n-1}^{(k)}} \prod_{1 \leq l < k} \mathbf{1}_{X_n^{(l)} = X_{n-1}^{(l)}}, & \text{if } X_n^{(r)} > X_{n-1}^{(r)}, \end{cases} \quad (2.6)$$

so if there is no jump in the r th order maximum process we sample from the conditional distribution and if there is a jump, we note the new value that caused the jump. For the second sequence we have $\widetilde{X}_{\uparrow+1,n} := -\infty$ if $n \leq r$ and if $n > r$

$$\widetilde{X}_{\uparrow+1,n} := \begin{cases} F^{\leftarrow}(U_{r,n}|X_n^{(r)}) & \text{if } X_n^{(r)} = X_{n-1}^{(r)}, \\ X_{n-1}^{(r)} & \text{if } X_n^{(r)} > X_{n-1}^{(r)}, \end{cases} \quad (2.7)$$

so if there is no jump in the r th order maximum at n , we sample from the conditional distribution and if there is a jump at index n we note the smaller value at $n - 1$ that the process jumps from. The sequence $\widetilde{X}_{\uparrow+1}$ depends on $X, X^{(1)}, \dots, X^{(r)}$ only via $X^{(r)}$, but $\widehat{X}_{\uparrow+1}$ depends on all $X^{(1)}, X^{(2)}, \dots, X^{(r)}$.

2.3 Identities in law and the Markov property

Next we provide some identities in law which will show that the sequence $X^{(r)}, r \geq 1$ of extremal processes is a sequence-valued Markov chain.

Theorem 2.1 *For $r \in \mathbb{N}$ the following random variables are equal in distribution as random elements in $(\mathbb{R}_{-\infty}^{\mathbb{N}})^{(r+1)}$,*

$$(X^{(0)}, X^{(1)}, \dots, X^{(r)}) \stackrel{d}{=} (\widehat{X}_{\uparrow+1}, X^{(1)}, \dots, X^{(r)}), \quad (2.8)$$

and

$$(X^{(1)}, \dots, X^{(r+1)}) \stackrel{d}{=} (X^{(1)}, \dots, X^{(r)}, \widetilde{X}_{\uparrow+1}). \quad (2.9)$$

In particular, $X^{(1)}, X^{(2)}, \dots$ is a Markov chain with state space $\mathbb{R}_{-\infty}^{\mathbb{N}, \uparrow}$, with its conditional distributions satisfying

$$X^{(r+1)} \Big| X^{(r)}, \dots, X^{(1)} \stackrel{d}{=} \widetilde{X}_{\uparrow+1} \Big| X^{(r)}, \quad r \in \mathbb{N}. \quad (2.10)$$

Proof Indeed, Eq. 2.9 follows from Eq. 2.8 because

$$\begin{aligned} (X^{(1)}, \dots, X^{(r+1)}) &= (X^{(1)}, \dots, X^{(r)}, (X^{(r)} \mathcal{R} \wedge X^{(0)})) \quad (\text{by Proposition 2.1}), \\ &\stackrel{d}{=} (X^{(1)}, \dots, X^{(r)}, (X^{(r)} \mathcal{R} \wedge \widehat{X}_{\uparrow+1})) \quad (\text{from (2.8)}) \\ &= (X^{(1)}, \dots, X^{(r)}, \widetilde{X}_{\uparrow+1}). \end{aligned}$$

The last equality holds because $\bigvee (X^{(r)} \mathcal{R} \wedge \widehat{X}_{(r+1)})$ has (for its finite components, when $n \geq r+1$) the terms $\bigvee_{j=r+1}^n (X_{j-1}^{(r)} \wedge \widehat{X}_{(r+1),j})$ (compare with Eq. 2.2), and in this, by Eq. 2.6, we take $\widehat{X}_{(r+1),j} = F^{\leftarrow}(U_{r,j}|X_j^{(r)})$ if $X_j^{(k)} = X_{j-1}^{(k)}$, $1 \leq k \leq r$; otherwise, there is a k , $1 \leq k \leq r$, with $X_j^{(k)} > X_{j-1}^{(k)}$ and $X_j^{(\ell)} = X_{j-1}^{(\ell)}$ for $1 \leq \ell < k$, in which case, by Eq. 2.6, we take $\widehat{X}_{(r+1),j} = X_j^{(r)}$. In the first case, $X_{j-1}^{(r)} \wedge \widehat{X}_{(r+1),j} = X_{j-1}^{(1)} \wedge \widehat{X}_{(r+1),j} = \widehat{X}_{(r+1),j} = F^{\leftarrow}(U_{r,j}|X_j^{(r)})$, and in the second case, $X_{j-1}^{(r)} \wedge \widehat{X}_{(r+1),j} = X_{j-1}^{(r)} \wedge X_j^{(r)} = X_{j-1}^{(r)}$. On taking $\bigvee_{j=r+1}^n$, this replicates the corresponding component for $\bigvee \widehat{X}_{(r+1)}$ (see Eq. 2.7).

Thus indeed (2.9) holds, and on the righthand side $\widehat{X}_{(r+1)}$ depends on $X^{(1)}, \dots, X^{(r)}$ only through $X^{(r)}$, $r \in \mathbb{N}$. In particular, Eq. 2.10 holds, and $X^{(1)}, X^{(2)}, \dots$ must be a Markov chain.

It remains to show (2.8). For $r \in \mathbb{N}$ let

$$\mathbb{R}_{-\infty}^{r,\downarrow} := \{\mathbf{m} = (m_1, \dots, m_r) \in \mathbb{R}_{-\infty}^r : m_1 \geq \dots \geq m_r\}$$

be the space of r -tuples with nonincreasing $\mathbb{R}_{-\infty}$ -valued components, and introduce a continuous truncation mapping $\mu_r = (\mu_{r,1}, \dots, \mu_{r,r}) : \mathbb{R}_{-\infty}^{r,\downarrow} \times \mathbb{R} \rightarrow \mathbb{R}_{-\infty}^{r,\downarrow}$, by setting

$$\mu_{r,1}(\mathbf{m}, x) = x \vee m_1,$$

and

$$\mu_{r,k}(\mathbf{m}, x) = m_{k-1} \mathbf{1}_{x > m_{k-1}} + m_k \mathbf{1}_{x \leq m_k} + x \mathbf{1}_{m_k < x \leq m_{k-1}}, \quad 2 \leq k \leq r, \quad (2.11)$$

when $x \in \mathbb{R}$ and $\mathbf{m} = (m_1, \dots, m_r) \in \mathbb{R}_{-\infty}^{r,\downarrow}$. Note that $\mu_{r,k}(\mathbf{m}, x)$ interpolates continuously between components m_k and m_{k-1} , $m_k \leq m_{k-1}$, of \mathbf{m} , and satisfies

$$\mu_{r,k}(\mathbf{m}, x) \geq m_k \text{ for } \mathbf{m} \in \mathbb{R}_{-\infty}^{r,\downarrow}, x \in \mathbb{R}, 1 \leq k \leq r. \quad (2.12)$$

Having constructed μ_r , define two further mappings

$$\widetilde{\mu}_r = (\widetilde{\mu}_{r,0}, \dots, \widetilde{\mu}_{r,r}) : \mathbb{R}_{-\infty}^{r,\downarrow} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}_{-\infty}^{r,\downarrow}$$

and

$$\widehat{\mu}_r = (\widehat{\mu}_{r,0}, \dots, \widehat{\mu}_{r,r}) : \mathbb{R}_{-\infty}^{r,\downarrow} \times \mathbb{R} \times (0, 1) \rightarrow \mathbb{R} \times \mathbb{R}_{-\infty}^{r,\downarrow},$$

as follows. Take $\mathbf{m} = (m_1, \dots, m_r) \in \mathbb{R}_{-\infty}^{r,\downarrow}$, $x \in \mathbb{R}$ and $u \in (0, 1)$. When $k = 0$, set

$$\widetilde{\mu}_{r,0}(\mathbf{m}, x) = x \quad \text{and} \quad \widehat{\mu}_{r,0}(\mathbf{m}, x, u) = \begin{cases} F^{\leftarrow}(u|m_r), & \text{if } x \leq m_r, \\ x, & \text{if } x > m_r. \end{cases} \quad (2.13)$$

When $1 \leq k \leq r$, set

$$\widetilde{\mu}_{r,k}(\mathbf{m}, x) = \mu_{r,k}(\mathbf{m}, x) \quad \text{and also} \quad \widehat{\mu}_{r,k}(\mathbf{m}, x, u) = \mu_{r,k}(\mathbf{m}, x). \quad (2.14)$$

With these mappings the component form of the lefthand side of Eq. 2.8 can be written as

$$(X_n, X_n^{(1)}, \dots, X_n^{(r)}, n \geq 2) = \widetilde{\mu}_r(X_{n-1}^{(1)}, \dots, X_{n-1}^{(r)}, X_n), \quad n \geq 2) \quad (2.15)$$

and the component form of the righthand side of Eq. 2.8 can be written as

$$(\widehat{X}_{(r+1),n}, X_n^{(1)}, \dots, X_n^{(r)}, n \geq 2) = \widehat{\mu}_r(X_{n-1}^{(1)}, \dots, X_{n-1}^{(r)}, X_n, U_{r,n}), n \geq 2). \quad (2.16)$$

We check that these are verified, as follows. Apply the formulae Eqs. 2.13 and 2.14, substituting $\mathbf{m} = (X_n^{(1)}, \dots, X_n^{(r)})$, $x = X_n$ and $u = U_{r,n}$. Consider the righthand side of Eq. 2.15. With those substitutions, the $k = 0$ component equals $\tilde{\mu}_{r,0}(\mathbf{m}, x) = x = X_n$, matching the lefthand side of Eq. 2.15. The k th component, for $1 \leq k \leq r$, with the substitutions, equals, by Eq. 2.11,

$$\begin{aligned} \tilde{\mu}_{r,k}(\mathbf{m}, x) &= \mu_{r,k}(\mathbf{m}, x) = X_{n-1}^{(k-1)} \mathbf{1}_{X_n > X_{n-1}^{(k-1)}} + X_{n-1}^{(k)} \mathbf{1}_{X_n \leq X_{n-1}^{(k)}} \\ &\quad + X_n \mathbf{1}_{X_{n-1}^{(k)} < X_n \leq X_{n-1}^{(k-1)}} = X_n^{(k)}, \end{aligned} \quad (2.17)$$

again matching the k th component on the lefthand side of Eq. 2.15. Next consider the righthand side of Eq. 2.16. With the above substitutions, the $k = 0$ component equals, by Eq. 2.13,

$$\widehat{\mu}_{r,0}(\mathbf{m}, x, u) = \begin{cases} F^{\leftarrow}(U_{r,n} | X_{n-1}^{(r)}), & \text{if } X_n \leq X_{n-1}^{(r)}, \\ X_n, & \text{if } X_n > X_{n-1}^{(r)}, \end{cases}$$

agreeing with $\widehat{X}_{(r+1),n}$ from Eq. 2.6. So the righthand side of Eq. 2.16 matches the lefthand side of Eq. 2.16 for the $k = 0$ component. The k th component, for $1 \leq k \leq r$, with the substitutions, equals, by Eq. 2.14, the righthand side of Eq. 2.17. So the righthand side of Eq. 2.16 matches the lefthand side of Eq. 2.16 for the components $1 \leq k \leq r$. With these checkings we have verified Eqs. 2.15 and 2.16.

In Eqs. 2.15 and 2.16, $X_n \perp (X_{n-1}^{(1)}, \dots, X_{n-1}^{(r)})$ and $U_{r,n} \perp (X_{n-1}^{(1)}, \dots, X_{n-1}^{(r)}, X_n)$ since we assumed that \mathbf{X} and \mathbf{U} are independent arrays of iid rv's. The right sides of Eqs. 2.15 and 2.16 are Markov chains with stationary transition probabilities in the index n (new value is a function of the previous value and an independent quantity) and for $n = 1$, the left sides of Eqs. 2.15 and 2.16 have common initial value $(X_1, X_1, -\infty, \dots, -\infty) \in \mathbb{R} \times \mathbb{R}_{-\infty}^{r,\downarrow}$. Therefore, to prove equality in distribution in Eq. 2.8, it suffices to prove both chains have a common transition kernel.

To see this, let $X' \stackrel{d}{=} X_1 \sim F$ and $U' \stackrel{d}{=} U_{1,1} \in (0, 1)$ be independent rv's. For $x, y \in \mathbb{R}$ with $F(y) > 0$ note

$$P(X' \leq y, F^{\leftarrow}(U' | y) \leq x) = P(X' \leq y)P(X' \leq x | X' \leq y) = F(x \wedge y). \quad (2.18)$$

Take $\mathbf{m} = (m_1, \dots, m_r)$, $\mathbf{m}' = (m'_1, \dots, m'_r) \in \mathbb{R}_{-\infty}^{r,\downarrow}$ with $F(m'_k) > 0$ for $1 \leq k \leq r$, and $m'_0 := \infty$. By Eq. 2.16 we have for the transition probability,

$$\begin{aligned} P \left(\widehat{X}_{(r+1),n+1}, X_{n+1}^{(1)}, \dots, X_{n+1}^{(r)} \in (-\infty, x] \times \prod_{k=1}^r [-\infty, m_k] \right. \\ \left. \middle| \widehat{X}_{(r+1),n} = y, (X_n^{(1)}, \dots, X_n^{(r)}) = \mathbf{m}' \right) \\ = P \left(\widehat{\mu}_r(\mathbf{m}', X', U') \in (-\infty, x] \times \prod_{k=1}^r [-\infty, m_k] \right). \end{aligned}$$

Decompose the last expression as

$$\begin{aligned} & P \left[\widehat{\mu}_r(\mathbf{m}', X', U') \in (-\infty, x] \times \prod_{k=1}^r [-\infty, m_k], X' \leq m'_r \right] \\ & + \sum_{k=1}^r P \left[\widehat{\mu}_r(\mathbf{m}', X', U') \in (-\infty, x] \times \prod_{l=1}^r [-\infty, m_l], X' \in (m'_k, m'_{k-1}] \right) \\ & =: A + B. \end{aligned}$$

First consider the probability A . It equals

$$\begin{aligned} & P \left[F^{\leftarrow}(U' | m'_r) \leq x, \mu_{rk}(\mathbf{m}', X', U') \leq m_k, 1 \leq k \leq r; X' \leq m'_r \right] \\ & = P \left[X' \leq x, \mu_{rk}(\mathbf{m}', X', U') \leq m_k, 1 \leq k \leq r; X' \leq m'_r \right). \end{aligned} \quad (2.19)$$

Next consider the probability B . For this we use Eq. 2.13 and get

$$B = \sum_{k=1}^r P(X' \leq x, \mu_{rl}(\mathbf{m}', X') \leq m_l, 1 \leq l \leq r, X' \in (m'_k, m'_{k-1}]), \quad (2.20)$$

in which $\mu_{rl}(\mathbf{m}', X') \geq m'_l, 1 \leq l \leq r$, by Eq. 2.12.

On the other hand, from the left sides of Eqs. 2.8 and 2.15,

$$\begin{aligned} & P \left((X_{n+1}, X_{n+1}^{(1)}, \dots, X_{n+1}^{(r)} \in (-\infty, x] \times \prod_{k=1}^r [-\infty, m_k] \right. \\ & \left. \middle| X_n = y, (X_n, X_n^{(1)}, \dots, X_n^{(r)}) = \mathbf{m}' \right) \\ & = P(\tilde{\mu}_r(\mathbf{m}', X') \in (-\infty, x] \times \prod_{k=1}^r [-\infty, m_k]) \\ & = P \left(X', \mu_{rl}(\mathbf{m}', X'), l = 1, \dots, r \in (-\infty, x] \times \prod_{k=1}^r [-\infty, m_k] \right). \end{aligned}$$

Again decompose the last as

$$\begin{aligned} & P \left(X' \leq x, \mu_{rl}(\mathbf{m}', X') \leq m_l, l = 1, \dots, r, X' \leq m'_r \right. \\ & \left. + \sum_{k=1}^r P(X' \leq x, \mu_{rl}(\mathbf{m}', X') \leq m_l, l = 1, \dots, r, X' \in (m_k, m_{k-1}]) \right) \\ & = A + B \quad (\text{by Eqs. 2.19 and 2.20}). \end{aligned}$$

This completes the proof of Eq. 2.8 and of Theorem 2.1. \square

Remark Probabilities A and B can be calculated explicitly as follows.

For A , take $m'_k > m_k$ for some $k = 1, \dots, r$. Then because of Eq. 2.12, the probability A is 0. So assume that $m'_k \leq m_k$ for $1 \leq k \leq r$. Then the condition $X' \leq$

m'_r in A implies $X' \leq m'_k \leq m_k$, hence $\mu_{r,k}(\mathbf{m}', X', U') \leq m_k$ for $k = 1, \dots, r$, by Eq. 2.11. So, using (2.18), A reduces to

$$A = P(F^{\leftarrow}(U'|m'_r) \leq x, X' \leq m'_r) = F(x \wedge m'_r) \prod_{1 \leq k \leq r} \mathbf{1}_{m'_k \leq m_k}. \quad (2.21)$$

For B , fix k and suppose $l > k$. Then the interval $(m'_l, m'_{l-1}]$ is to the left of $(m'_k, m'_{k-1}]$ where X' is located, and $\mu_{r,l}(\mathbf{m}', X') = m'_{l-1}$. The probability is then 0 unless $m_l \geq m'_{l-1}$. If $l < k$, the order of the intervals is reversed, $\mu_{r,l}(\mathbf{m}', X') = m'_l$, and the probability is 0 unless $m'_l \leq m_l$. Thus, B becomes

$$B = \sum_{k=1}^r \left[P(m'_k < X' \leq x \wedge m_k \wedge m'_{k-1}) \prod_{1 \leq l < k} \mathbf{1}_{m'_l \leq m_l} \prod_{k < l \leq r} \mathbf{1}_{m'_{l-1} \leq m_l} \right]. \quad (2.22)$$

3 Asymptotic behavior of the discrete time process $M^{(r)}$ for large r

In this section we consider the asymptotic behavior as $r \rightarrow \infty$ of the \mathbb{R}^∞ -valued stochastic process $\{\mathbf{M}^{(r)} := (M_n^{(r)}, n \geq r), r \geq 1\}$. As r increases the sequence moves further and further from its largest values, so limit behavior for both the range of $\mathbf{M}^{(r)}$ and $\mathbf{M}^{(r)}$ itself, depend critically on left tail behavior of the distribution of X_1 . Appropriate left-tail conditions related to minimal domains of attraction in classical extreme value theory make the range and the sequence of r th order maxima converge weakly.

Throughout Section 3, the underlying distribution F of the iid sequence $\{X_n\}$ is continuous, so the records are Poisson with atomless mean measure $R(\cdot)$ which has distribution $R(x) = -\log(1 - F(x))$ (Resnick 2008, page 166). The assumption of continuity could be relaxed as in (Engelen et al. 1988; Shorrock 1974; 1975) but results are most striking and elegant when F is continuous and we proceed in this setting. We assume $F(x)$ has left endpoint ℓ_F and right endpoint u_F so that the measure F has support $[\ell_F, u_F] \subset [-\infty, \infty]$.

3.1 r^{th} maximum and r -records

Assume $F(x) < 1$ and define

$$\begin{aligned} R_n &= \sum_{j=1}^n \mathbf{1}_{[X_j \geq X_n]} = \text{relative rank of } X_n \text{ among } X_1, \dots, X_n \\ &= \text{rank of } X_n \text{ at "birth"}. \end{aligned}$$

The $\{R_n\}$ are independent random variables and R_n is uniformly distributed (Rényi 1962) on $\{1, \dots, n\}$; that is,

$$P(R_n = i) = 1/n, \quad i = 1, \dots, n.$$

Considering $\{\mathbf{M}^{(r)}, r \geq 1\}$ as an \mathbb{R}^∞ -valued stochastic process, we ask for the asymptotic behavior of $\mathbf{M}^{(r)}$ and its range as a function of r as $r \rightarrow \infty$.

Define the r -record times of $\{X_n\}$ by

$$L_0^{(r)} = 0, \quad L_{n+1}^{(r)} = \inf\{j > L_n^{(r)} : R_j = r\}.$$

The r -records are $\{X_{L_n^{(r)}}, n \geq 1\}$. Ignatov's theorem (Ignatov 1986; Goldie and Rogers 1984; Engelen et al. 1988; Resnick 2008) says that

$$\{\{X_{L_n^{(r)}}, n \geq 1\}, r \geq 1\}$$

are iid Poisson processes, each with mean measure $R(\cdot)$ on \mathbb{R} . A Poisson process with mean measure R is denoted $\text{PRM}(R)$.

We list some initial facts about $\mathbf{M}^{(r)}$ and its range.

- For fixed r , $\mathbf{M}^{(r)} = \{M_n^{(r)}, n \geq r\}$ jumps at time $k \geq r$ iff

$$R_k \in \{1, \dots, r\}.$$

So the events

$$\{\mathbf{M}^{(r)} \text{ jumps at time } k, k \geq r\}$$

are independent events over k and

$$P(\mathbf{M}^{(r)} \text{ jumps at } k) = \frac{r}{k}.$$

Remark 3.1 This has the implication that if we re-index and set $k = r + l$ for $l \geq 0$, then for any fixed l , as $r \rightarrow \infty$,

$$P(\mathbf{M}^{(r)} \text{ jumps at } r + l) = \frac{r}{r + l} \rightarrow 1.$$

So for large r , $\mathbf{M}^{(r)}$ jumps at almost every integer. Define the jump indices

$$\{\tau_l^{(r)}, l \geq 0\} = \{j \geq 1 : M_{r+j}^{(r)} > M_{r+j-1}^{(r)}\} \cup \{0\}.$$

Then in \mathbb{R}_+^∞ ,

$$\{\tau_l^{(r)}, l \geq 0\} \Rightarrow \{0, 1, 2, \dots\}.$$

- For fixed r , let \mathcal{R}_r be the range of $\mathbf{M}^{(r)}$; that is, the distinct points without repetition in the sequence $\{M_n^{(r)}, n \geq r\}$. See Fig. 1. Then,

$$\mathcal{R}_r := \bigcup_{p=1}^r \bigwedge \{X_{L_n^{(p)}}, n \geq 1\}, \quad (3.1)$$

By Ignatov's theorem (Ignatov 1986; Stam 1985; Goldie and Rogers 1984; Engelen et al. 1988; Resnick 2008), this is a sum of r independent $\text{PRM}(R)$ processes and therefore the range of $\mathbf{M}^{(r)}$ is $\text{PRM}(rR)$.

To prove (3.1), suppose $M_n^{(r)} = x$ for some $n \geq r$. Suppose the r th largest of X_1, \dots, X_n occurs at $X_i = x$ for $i \leq n$. If the rank of X_i were $> r$, it could not be the case that $M_n^{(r)} = x$. This shows that

$$\text{range of } \mathbf{M}^{(r)} \subset \bigcup_{p=1}^r \bigwedge \{X_{L_n^{(p)}}, n \geq 1\}.$$

Conversely, suppose $X_{L_n^{(p)}} = x$, so at time $L_n^{(p)}$, the rank of $X_{L_n^{(p)}}$ is p . Wait until $r - p$ additional X 's have been observed that exceed x and then the r th largest will equal x .

3.2 Limits for the range \mathcal{R}_r of $M^{(r)}$

Although our primary interest is in the behavior of $\{M^{(r)}, r \geq 1\}$ as an \mathbb{R}^∞ -valued random sequence, it is instructive and helpful to discuss the behavior of the range \mathcal{R}_r of $M^{(r)}$.

As a basic result we derive a deterministic limit for \mathcal{R}_r . Let \mathcal{R} be the support of the measure $R(\cdot)$ which is also the support of F .

Proposition 3.2 *As $r \rightarrow \infty$, \mathcal{R}_r , the range of $M^{(r)}$, converges as a random closed set in the Fell topology (Molchanov 2005; Matheron 1975; Vervaat and Holwerda 1997) on $[\ell_F, u_F]$ to the non-random limit \mathcal{R} :*

$$\mathcal{R}_r \Rightarrow \mathcal{R}. \quad (3.2)$$

Proof Since $\mathcal{R}_r \subset \mathcal{R}$, it suffices to show for any open G with $\mathcal{R} \cap G \neq \emptyset$, that

$$P(\mathcal{R}_r \cap G \neq \emptyset) \rightarrow 1.$$

However, $\mathcal{R} \cap G \neq \emptyset$ implies $R(G) > 0$ and therefore,

$$\begin{aligned} P(\mathcal{R}_r \cap G \neq \emptyset) &= 1 - P(\text{PRM}(rR(G)) = 0) \\ &= 1 - e^{-rR(G)} \rightarrow 1, \quad (r \rightarrow \infty) \end{aligned}$$

since $R(G) > 0$. □

The set convergence in Eq. 3.2 is to a deterministic limit. Since \mathcal{R}_r is a $\text{PRM}(rR)$ point process, we can get a random limit if we center and scale the $\{X_n\}$ so that the mean measure rR converges to a Radon measure. Recall $R(x) = -\log \bar{F}(x)$.

Assume there exist $a_r > 0$ and $b_r \in \mathbb{R}$ and a non-decreasing limit function $g(x)$ with more than one point of increase such that

$$rR(a_r x - b_r) \rightarrow g(x), \quad (r \rightarrow \infty). \quad (3.3)$$

For x such that $g(x) > 0$, we must have $R(a_r x - b_r) \rightarrow 0$ and thus $a_r x - b_r$ converging to the left endpoint of F (and R) to counteract $r \rightarrow \infty$. We now explain why e^{-g} is related to an extreme value distribution. Remembering that $e^{-R} = \bar{F}$, Eq. 3.3 is equivalent to

$$(\bar{F}(a_r x - b_r))^r = \exp\{-rR(a_r x - b_r)\} \rightarrow e^{-g(x)}$$

or

$$P \left(\frac{\wedge_{i=1}^r X_i + b_r}{a_r} > x \right) \rightarrow e^{-g(x)}. \quad (3.4)$$

So we recognize e^{-g} as the survivor function of an extreme value distribution of minima of iid random variables. Expressing this in terms of maxima by setting $Y_i = -X_i$ we get Eq. 3.4 equivalent to

$$P \frac{\bigvee_{i=1}^r Y_i - b_r}{a_r} \leq -x \rightarrow e^{-g(x)} = G_\gamma(-x), \quad (3.5)$$

for some $\gamma \in \mathbb{R}$, where $G_\gamma(x) = \exp\{-(1 + \gamma x)^{-1/\gamma}\}$, $1 + \gamma x > 0$ is the shape parameter family of extreme value distributions for maxima (Resnick 2008; de Haan and Ferreira 2006). So in Eq. 3.3, $g(x) = g_\gamma(x) = -\log G_\gamma(-x)$. The equivalent way to write (3.5) is

$$rP(Y_1 > a_r(-x) + b_r \rightarrow g(x), \quad \forall x \text{ s.t. } g(x) > 0,$$

and Eq. 3.3 is equivalent to

$$rF(a_r x - b_r) \rightarrow g(x), \quad \forall x \text{ s.t. } g(x) > 0. \quad (3.6)$$

In particular, apart from centering, we have the cases:

(1) *Gumbel case*: $\gamma = 0$. Then

$$g_0(x) = e^x, \quad x \in \mathbb{R}.$$

(2) *Reverse Weibull case*: $\gamma < 0$: Then $1 + \gamma(-x) > 0$ iff $x > -1/|\gamma|$ and

$$g_\gamma(x) = (1 + |\gamma|x)^{1/|\gamma|}, \quad x > -1/|\gamma|.$$

Adjusting the centering and scaling by taking $b_r = 0$, we find R is regularly varying at 0 and

$$rR(a_r x) \rightarrow x^{1/|\gamma|}, \quad x > 0.$$

(3) *Frechét case*: $\gamma > 0$. Then $1 + \gamma(-x) > 0$ iff $x < 1/\gamma$ and

$$g_\gamma(x) = (1 - \gamma x)^{-1/\gamma}, \quad x < 1/\gamma.$$

Adjusting the centering and scaling so the support is $(-\infty, 0)$, we get

$$rR(a_r x) \rightarrow |x|^{-1/\gamma}, \quad x < 0,$$

which implies regular variation at 0 from the left.

We can apply this analysis to get convergence of \mathcal{R}_r after centering and scaling. Recall \mathcal{R}_r is PRM(rR). A family of Poisson point measures converges weakly iff the mean measures converge (eg. Resnick (2007)). So replacing

$$X_i \rightarrow \frac{X_i + b_r}{a_r}$$

rescales the points of the range to be Poisson with mean measure given by the left side of Eq. 3.3. Let

$$\text{supp}_\gamma = \{x : 1 - \gamma x > 0\} \quad (3.7)$$

and $m_\gamma(\cdot)$ be the measure with density $g'_\gamma(x)$, $x \in \text{supp}_\gamma$. Let $M_+(\text{supp}_\gamma)$ be the space of Radon measures on supp_γ , topologized by vague convergence. Then Eq. 3.3 implies the vague convergence

$$rR\left(\frac{\cdot - b_r}{a_r}\right) \xrightarrow{v} m_\gamma(\cdot)$$

in $M_+(\text{supp}_\gamma)$, and thus on $M_+(\text{supp}_\gamma)$ we have

$$(\mathcal{R}_r + b_r)/a_r \Rightarrow \text{PRM}(m_\gamma). \quad (3.8)$$

We may realize $\text{PRM}(m_\gamma)$ as follows: Let $\Gamma_i = \sum_{j=1}^i E_j$ be a sum of iid standard exponential random variables. The $\{\Gamma_i\}$ are points of a homogeneous Poisson process rate 1 on $[0, \infty)$. The measure m_γ has distribution

$$g_\gamma : \text{supp}_\gamma \rightarrow (0, \infty),$$

with inverse

$$g_\gamma^\leftarrow : (0, \infty) \rightarrow \text{supp}_\gamma.$$

The transformation theory for Poisson processes (Resnick 2007, Section 5.1)) means that if we map homogeneous Poisson points $\{\Gamma_i, i \geq 1\}$ to $\{g_\gamma^\leftarrow(\Gamma_i), i \geq 1\}$, these become the points of $\text{PRM}(m_\gamma)$ on supp_γ . For instance, if $\gamma = 0$, $\text{supp}_0 = \mathbb{R}$, $g_0(x) = e^x$, $x \in \mathbb{R}$, and $g_0^\leftarrow(y) = \log y$, $y > 0$, then $\text{PRM}(m_0)$ has points $\{\log \Gamma_i, i \geq 1\}$.

3.3 Weak convergence of the r th maxima sequence $M^{(r)}$

Having understood how to get the range \mathcal{R}_r of $M^{(r)}$ to converge, we turn to convergence of $M^{(r)}$ itself. We continue to suppose the minimum domain of attraction condition, so that R satisfies (3.3), and recall $M_+(\text{supp}_\gamma)$ is the space of Radon measures on supp_γ , topologized by vague convergence. Point measures in $M_+(\text{supp}_\gamma)$ are denoted by $\sum_i \epsilon_{x_i}(\cdot)$ where $\epsilon_x(\cdot)$ is the Dirac measure placing mass 1 at x .

We start with a preliminary result on the empirical measures generated by $\{X_i\}$ that will be needed to study the weak convergence of $\{M^{(r)}\}$.

Proposition 3.3 *Assume (3.3). If N is a random element of $M_+(\text{supp}_\gamma)$ which is $\text{PRM}(m_\gamma)$, then for any $j \geq 0$,*

$$\sum_{i=1}^{r+j} \epsilon_{(X_i+b_r)/a_r} \Rightarrow N = \sum_{i=1}^{\infty} \epsilon_{g_\gamma^\leftarrow(\Gamma_i)} = \text{PRM}(m_\gamma), \quad (3.9)$$

in $M_+(\text{supp}_\gamma)$ and, in fact, jointly for any $k \geq 0$,

$$\sum_{i=1}^{r+j} \epsilon_{(X_i+b_r)/a_r; 0 \leq j \leq k} \Rightarrow (N, \dots, N) \quad (3.10)$$

in $M_+(\text{supp}_\gamma) \times \dots \times M_+(\text{supp}_\gamma)$.

Proof We have Eq. 3.10 following from Eq. 3.9 since with respect to the vague distance $d(\cdot, \cdot)$ on $M_+(\text{supp}_\gamma)$ (see, eg. (Resnick 2007, page 51))

$$d\left(\sum_{i=1}^r \epsilon_{(X_i+b_r)/a_r}, \sum_{i=1}^{r+j} \epsilon_{(X_i+b_r)/a_r}\right) \Rightarrow 0$$

for any $j > 0$. To verify this, let f be positive and continuous with compact support on supp_γ and from Eq. (3.14) of (Resnick 2007, p.51), it suffices to show

$$E \left| \sum_{i=1}^r f((X_i + b_r)/a_r) - \sum_{i=1}^{r+j} f((X_i + b_r)/a_r) \right| \rightarrow 0.$$

The difference is

$$E \sum_{i=r+1}^{r+j} f((X_i + b_r)/a_r) = E \sum_{i=1}^j f((X_i + b_r)/a_r)$$

and assuming the support of f is a compact set K in supp_γ , this is bounded above by

$$\sup_{x \geq 0} f(x) j P[X_1 \in a_r K - b_r] \rightarrow 0,$$

since for $x \in K$, $a_r x - b_r$ converges to the left endpoint of F , and, under (3.3), there cannot be an atom at this left endpoint.

The result in Eq. 3.9 follows by a small modification of the proof of Theorem 5.3 in (Resnick 2007, p.138) since (3.3) is equivalent to Eq. 3.6. \square

Now we turn to \mathbb{R}^∞ -convergence of the r th maximum sequence. Continue to suppose (3.3). Without normalization, the sequence $\mathbf{M}^{(r)}$ converges to a sequence all of whose entries are the left endpoint of F . In order to get $\mathbf{M}^{(r)}$ to converge, we must have $M_r^{(r)} = \wedge_{i=1}^r X_i$ converge and this helps explain why a domain of attraction condition for minima is relevant. The condition (3.3) produces a non-trivial limit.

Proposition 3.4 *Suppose the domain of attraction condition (3.3) holds. Then in \mathbb{R}^∞ ,*

$$\frac{M_r^{(r)} + b_r}{a_r} = \frac{M_{r+j}^{(r)} + b_r}{a_r}, j \geq 0 \Rightarrow g_\gamma^\leftarrow(\Gamma_l, l \geq 1) \quad (r \rightarrow \infty), \quad (3.11)$$

where $\{\Gamma_l, l \geq 1\}$ are the points of a homogeneous Poisson process on \mathbb{R}_+ .

Proof Fix $j \geq 0$ and observe for $x \in \text{supp}_\gamma$,

$$\left\{ \frac{M_{r+j}^{(r)} + b_r}{a_r} > x \right\} = \left\{ \sum_{i=1}^{r+j} \epsilon_{(X_i + b_r)/a_r}(x, \infty) \geq r \right\} = \left\{ \sum_{i=1}^{r+j} \epsilon_{(X_i + b_r)/a_r}((-\infty, x]) \leq j \right\}$$

and therefore

$$\left\{ \frac{M_{r+j}^{(r)} + b_r}{a_r} \leq x \right\} = \left\{ \sum_{i=1}^{r+j} \epsilon_{(X_i + b_r)/a_r}((-\infty, x]) > j \right\}.$$

For a non-decreasing sequence $\{x_j\}$ of real numbers in supp_γ ,

$$P \bigcap_{j=0}^k \left[\frac{M_{r+j}^{(r)} + b_r}{a_r} \leq x_j \right] = P \bigcap_{j=0}^k \left[\sum_{i=1}^{r+j} \epsilon_{(X_i + b_r)/a_r}([-\infty, x_j]) > j \right]$$

and applying (3.10) yields that the RHS converges to

$$\begin{aligned} P \left(\bigcap_{j=0}^k [N((-\infty, x_j] > j) \right] &= P \sum_{i=1}^{\infty} \epsilon_{g_Y^{\leftarrow}(\Gamma_i)}(-\infty, x_j] > j; j = 0, \dots, k) \\ &= P \left(g_Y^{\leftarrow}(\Gamma_{j+1}) \leq x_j; j = 0, \dots, k \right). \end{aligned}$$

This yields the announced result (3.11). \square

To summarize: Without normalization, the random set consisting of the range of $M^{(r)}$ converges to the deterministic limit consisting of the support of F . To get a random limit requires the minimum domain of attraction condition and then the centered and scaled range converges to a limit Poisson process. Likewise, convergence in distribution of the \mathbb{R}^∞ -valued random elements $M^{(r)}$ as $r \rightarrow \infty$ requires the minimal domain of attraction condition.

4 Continuous time r th-order extremal processes

This section transitions to continuous time problems. The treatment is parallel to what we gave for discretely indexed processes but here the processes are generated by two-dimensional Poisson processes on $\mathbb{R}_+ \times \mathbb{R}$ and correspond to r th order extremal processes. One example of an r th order extremal process is obtained by taking the r th largest jump of a Lévy process up to time $t > 0$.

The continuous time case differs from the discrete index case, in that there are always infinitely many values *below* your present position. This necessitates differences in treatment. In continuous time we obtain modifications of Brownian motion limits whereas in discrete time we obtain Poisson limits for the r th order extremes.

The setup is as follows. Given an infinite measure $\Pi(\cdot)$ on an interval (ℓ_Π, u_Π) satisfying $-\infty \leq \ell_\Pi < u_\Pi \leq \infty$, $\Pi(\ell_\Pi, u_\Pi) = \infty$ and $Q(x) := \Pi(x, u_\Pi) < \infty$ for $\ell_\Pi < x < u_\Pi$. Let

$$N = \sum_k \epsilon_{(t_k, j_k)}, \quad (4.1)$$

be Poisson random measure on $\mathbb{R}_+ \times (\ell_\Pi, u_\Pi)$, with mean measure $Leb \times \Pi$, where $Leb(\cdot)$ is Lebesgue measure on \mathbb{R}_+ . Recall $\epsilon_{(t,x)}(\cdot)$ is a Dirac measure with mass 1 at the point (t, x) . Sometimes we write $(t_k, j_k) \in \text{supp}(N)$ to indicate the point (t_k, j_k) is charged by N . We assume ℓ_Π and u_Π are not atoms of Π and in fact, to make results most elegant we assume $\Pi(\cdot)$ is atomless. (Otherwise, results would be stated in terms of simplifications of point processes; see Engelen et al. (1988).) Our assumptions mean that

- (1) The function $Q(x)$ satisfies $Q(u_\Pi) = 0$ and $Q(\ell_\Pi) = \infty$ so $Q : (\ell_\Pi, u_\Pi) \rightarrow (0, \infty)$ and $Q(x)$ is non-increasing.
- (2) For any $t > 0$ and $u_\Pi \geq x > \ell_\Pi$: $N([0, t] \times (x, u_\Pi)) < \infty$ almost surely.
- (3) For any $t > 0$ and $u_\Pi \geq x > \ell_\Pi$: $N([0, t] \times (\ell_\Pi, x)) = \infty$ almost surely.

Traditionally, the (first-order) extremal process is defined by Resnick (2008), Deheuvels (1983), Deheuvels (1982), Dwass (1974), Dwass (1966), Dwass (1964),

Resnick (1975), Resnick and Rubinovitch (1973), Resnick (1974), Shorrock (1975), and Weissman (1975),

$$Y(t) = Y^{(1)}(t) = \underset{t_k \leq t}{j_k}, \quad 0 < t < \infty,$$

the largest j_k whose t_k coordinate is at or before time t . Alternatively we may write

$$Y(t) = \inf\{x > \ell_\Pi : N([0, t] \times (x, u_\Pi)) = 0\} = \inf\{x > \ell_\Pi : N([0, t] \times (x, u_\Pi)) < 1\}.$$

We develop the analogs of Propositions 3.2 and 3.4 as $r \rightarrow \infty$ for the continuous time r th order extremal process $Y^{(r)} := \{Y^{(r)}(t), 0 < t < \infty\}$ defined as,

$$Y^{(r)}(t) := \inf\{x > \ell_\Pi : N([0, t] \times (x, u_\Pi)) < r\}, \quad t > 0. \quad (4.2)$$

This means for $t > 0$, $u_\Pi \geq x > \ell_\Pi$,

$$\{Y^{(r)}(t) > x\} = \{N([0, t] \times (x, u_\Pi)) \geq r\},$$

and therefore,

$$\{Y^{(r)}(t) \leq x\} = \{N([0, t] \times (x, u_\Pi)) < r\}. \quad (4.3)$$

Alternative ways of considering $Y^{(r)}$ are in (Engelen et al. 1988).

What is the behavior of $\{Y^{(r)}, r \geq 1\}$, considered as a sequence of random elements of càdlàg space $D(\ell_\Pi, u_\Pi)$, as $r \rightarrow \infty$? Unlike in Section 3.3, here there are always infinitely many points below your current position and thus the left tail condition (3.6) used for $M^{(r)}$ must be different when considering $Y^{(r)}$. We analyze the range of $Y^{(r)}$ and for the weak limit behavior of $Y^{(r)}$, instead of relying on Poisson behavior, we rely on asymptotic normality.

4.1 The range \mathcal{R}_r of $Y^{(r)}$

Let \mathcal{R}_r be the unique values in the set $\{Y^{(r)}(t), t > 0\}$. As in the discrete time case (3.1), we have

$$\mathcal{R}_r = \bigcup_{p=1}^r \left\{ j_k : (t_k, j_k) \in \text{supp}(N), N([0, t_k] \times [j_k, u_\Pi)) = p \right\}. \quad (4.4)$$

To verify (4.4) suppose $x \in \mathcal{R}_r$. There exists $t > 0$ such that $Y^{(r)}(t) = x$, and therefore there exists $(t_k, x) \in \text{supp}(N)$ such that $t_k \leq t$. If $N([0, t_k] \times [x, u_\Pi)) > r$, then $Y^{(r)}(t) > x$, giving a contradiction. Thus x is in the right side of Eq. 4.4. Conversely, suppose j_k satisfies that there exists t_k such that $(t_k, j_k) \in \text{supp}(N)$ and $N([0, t_k] \times [j_k, u_\Pi)) = p$ for some $p \leq r$. Then there exists $t > t_k$ such that $N((t_k, t] \times [j_k, u_\Pi)) = r - p$ and thus $Y^{(r)}(t) = j_k$. Therefore, j_k belongs to the left side of Eq. 4.4. Note that the sets $(s, t] \times [j, u_\Pi)$ are all continuity sets of the intensity and therefore $t \rightarrow N((t_k, t] \times [j_k, u_\Pi))$ jumps by 1 with probability one.

When Π is atomless, the range of $Y(t) = Y^{(1)}(t)$ is known to be a Poisson process with mean measure determined by the monotone function $S(x) := -\log Q(x) = -\log \Pi(x, u_\Pi)$, $x > \ell_\Pi$. This is discussed, for example, in (Resnick 2008, page 183). In fact, from (Engelen et al. 1988, Theorem 6.2, page 234), the p -records of N are iid in p , and each sequence of p -records forms PRM(S). (A p -record of N

is a point j_k such that there exists t_k making $(t_k, j_k) \in \text{supp}(N)$ and $N([0, t_k] \times [j_k, u_\Pi)) = p$.) This and Eq. 4.4 allow us to conclude that \mathcal{R}_r is a Poisson process with mean measure $rS(\cdot)$. This achieves the continuous time analog of the discrete time discussion at the beginning of Section 3.2, and without any normalization we have

$$\mathcal{R}_r \Rightarrow \text{supp}(S), \quad (r \rightarrow \infty),$$

in the Fell topology of closed subsets of (ℓ_Π, u_Π) .

Paralleling the discrete time analysis, we proceed to obtain a non-degenerate limit for \mathcal{R}_r . We have to be more careful in the continuous case. The reason is that \mathcal{R}_r is PRM with mean measure $rS(\cdot)$ and S is Radon on (ℓ_Π, u_Π) , and it may allocate infinite mass to a neighborhood of both ℓ_Π and u_Π . Recall $S(x)$ satisfies $S(\ell_\Pi) = -\infty$ and $S(u_\Pi) = \infty$.

Assume without loss of generality that $\ell_\Pi < 0 < u_\Pi$. (If this is not the case, choose an arbitrary point between ℓ_Π and u_Π .) We make a treatment parallel to the discrete one by splitting the Poisson points of \mathcal{R}_r into those above 0 and those below. So write

$$\mathcal{R}_r = \mathcal{R}_r^+ \cup \mathcal{R}_r^-$$

where \mathcal{R}_r^+ are the positive Poisson points of \mathcal{R}_r and \mathcal{R}_r^- are the negative points of \mathcal{R}_r . The two Poisson processes \mathcal{R}_r^\pm are independent because their points are in disjoint regions. Define the two non-decreasing functions on \mathbb{R}_+ ,

$$S^+(x) = S(0, x] = S(x) - S(0), \quad 0 < x \leq u_\Pi \quad (4.5)$$

$$S^-(x) = S[-x, 0) = S(0) - S(-x), \quad 0 < x \leq -\ell_\Pi. \quad (4.6)$$

Assume there exist $a^\pm(t) > 0$, $b^\pm(t) \in \mathbb{R}$ and infinite Radon measures S_∞^\pm on \mathbb{R}_+ such that as $t \rightarrow \infty$,

$$tS^+(a^+(t)x - b^+(t)) \rightarrow S_\infty^+(x), \quad (4.7)$$

$$tS^-(a^-(t)x - b^-(t)) \rightarrow S_\infty^-(x). \quad (4.8)$$

The form of S_∞^\pm is determined by defining probability distribution tails $\bar{H}^\pm(x)$ by

$$\bar{H}^+(x) = e^{-S^+(x)}, \quad 0 < x < u_\Pi, \quad (4.9)$$

$$\bar{H}^-(x) = e^{-S^-(x)}, \quad 0 < x < -\ell_\Pi. \quad (4.10)$$

Note $\bar{H}^\pm(0) = e^{-S^\pm(0)} = e^{-0} = 1$ and $\bar{H}^+(u_\Pi) = e^{-S^+(u_\Pi)} = e^{-\infty} = 0$ and $\bar{H}^-(-\ell_\Pi) = 0$, similarly. Then, as in the discussion following (3.3), we find for $\gamma^\pm \in \mathbb{R}$ that

$$e^{-S^\pm(x)} = G_{\gamma^\pm}(-x),$$

where $G_\gamma(x)$ has a form given after (3.5). Note, if we want

$$a^+(t) = a^-(t) \quad \text{and} \quad b^+(t) = b^-(t)$$

up to convergence of types, we would need (Resnick 1971), $-\ell_\Pi = u_\Pi$ and

$$\bar{H}^+(x) \sim \bar{H}^-(x) \quad (x \rightarrow u_\Pi).$$

We now summarize.

Theorem 4.1 *The two Poisson processes \mathcal{R}_r^\pm are independent with $\mathcal{R}_r = \mathcal{R}_r^+ \cup \mathcal{R}_r^-$ where \mathcal{R}_r^+ has mean measure rS^+ on \mathbb{R}_+ and $-\mathcal{R}_r^-$ has mean measure rS^- on \mathbb{R}_+ so that \mathcal{R}_r^- are points on $(-\infty, 0)$. As $r \rightarrow \infty$, the range centered and scaled converges to a limiting Poisson process,*

$$\left(\frac{\mathcal{R}_r^+ + b^+(r)}{a^+(r)}, \frac{-\mathcal{R}_r^- + b^-(r)}{a^-(r)} \right) \Rightarrow (\mathcal{R}_\infty^+, -\mathcal{R}_\infty^-),$$

where the limits are independent Poisson processes on \mathbb{R}_+ with mean measures S_∞^\pm . So if Eqs. 4.7 and 4.8 hold, centering positive and negative range points appropriately leads to a limiting Poisson process such that positive points have mean measure $S_\infty^+(\cdot)$ and negative range points made positive by taking absolute values have mean measure $S_\infty^-(\cdot)$.

4.2 Finite dimensional convergence of $Y^{(r)}$ as random elements of $D(\ell_\Pi, u_\Pi)$

In this subsection, we give a left-tail condition on $\Pi(\cdot)$ guaranteeing finite dimensional convergence of $Y^{(r)}$ to a transformed Brownian motion.

Suppose there exist normalizing functions $a(r) > 0$, $b(r) \in \mathbb{R}$, and a non-decreasing limit function $h(x) \in \mathbb{R}$ with at least two points of increase such that for $a(r)x + b(r) \in (\ell_\Pi, u_\Pi)$,

$$\lim_{r \rightarrow \infty} \frac{r - Q(a(r)x + b(r))}{\sqrt{r}} = h(x). \quad (4.11)$$

Implications:

- (1) If we divide in Eq. 4.11 by r instead of \sqrt{r} , the limit will be 0 and therefore,

$$Q(a(r)x + b(r)) \sim r, \quad (r \rightarrow \infty). \quad (4.12)$$

Therefore, since $r \rightarrow \infty$, we must have that $Q(a(r)x + b(r)) \rightarrow \infty$ and $(\ell_\Pi, u_\Pi) \ni a(r)x + b(r) \rightarrow \ell_\Pi$.

- (2) For any $t > 0$,

$$\begin{aligned} \frac{r - tQ(a(r/t)x + b(r/t))}{\sqrt{r}} &= t \frac{r/t - Q(a(r/t)x + b(r/t))}{\sqrt{r/t}\sqrt{t}} \\ &\rightarrow \sqrt{t}h(x), \quad (r \rightarrow \infty). \end{aligned} \quad (4.13)$$

- (3) If we write $r - Q = (\sqrt{r} - \sqrt{Q})(\sqrt{r} + \sqrt{Q})$ and use Eq. 4.12, we get

$$\sqrt{r} - \sqrt{Q(a(r)x + b(r))} \rightarrow \frac{1}{2}h(x). \quad (4.14)$$

Remember that Q is non-increasing and define a probability distribution function $G(x)$ by $G(x) := \exp\{-\sqrt{Q(x)}\}$, so that G concentrates on (ℓ_Π, u_Π) . Then exponentiate in Eq. 4.14 to get

$$e^{\sqrt{r}} e^{-\sqrt{Q(a(r)x + b(r))}} \rightarrow e^{\frac{1}{2}h(x)}, \quad (r \rightarrow \infty)$$

or, after a change of variables $s = e^{\sqrt{r}}$,

$$sG\left(\left((\log s)^2\right)x + b((\log s)^2)\right) = se^{-\sqrt{Q(a((\log s)^2)x + b((\log s)^2))}} \rightarrow e^{\frac{1}{2}h(x)}, \quad (4.15)$$

as $s \rightarrow \infty$. So we conclude that $G(x) := e^{-\sqrt{Q(x)}}$ is in a domain of attraction of an extreme value distribution for minima. This technique is essentially the same as the one used to study limit laws for record values in Resnick (1973) or Resnick (2008).

- (4) Form of $h(x)$: As we saw following (3.6), if $\exp\{\frac{1}{2}h(x)\}$ plays the role of $g(x)$ then $h(x)$ must be of the form

$$e^{\frac{1}{2}h(x)} = -\log G_\gamma(-x),$$

where G_γ is an extreme value distribution for maxima of the form

$$G_\gamma(x) = \exp\{-(1 + \gamma x)^{-1/\gamma}\}, \quad \gamma \in \mathbb{R}, \quad 1 + \gamma x > 0.$$

So

$$\frac{1}{2}h(x) = \begin{cases} -\frac{1}{\gamma} \log(1 - \gamma x), & \text{if } \gamma \neq 0, \quad 1 - \gamma x > 0, \\ x, & \text{if } \gamma = 0, \quad x \in \mathbb{R}. \end{cases} \quad (4.16)$$

Observe that $h : \text{supp}_\gamma \rightarrow \mathbb{R}$ and $h^\leftarrow : \mathbb{R} \rightarrow \text{supp}_\gamma$. Recalling the definition of supp_γ from Eq. 3.7, we have

$$\text{supp}_\gamma = \{x \in \mathbb{R} : 1 - \gamma x > 0\} = \begin{cases} (-\frac{1}{|\gamma|}, \infty), & \text{if } \gamma < 0, \\ (-\infty, \frac{1}{|\gamma|}), & \text{if } \gamma > 0, \\ \mathbb{R}, & \text{if } \gamma = 0. \end{cases}$$

We apply these findings to obtain a marginal limit distribution for $Y^{(r)}(t)$ under the left tail condition. Assume (4.11). We show that, for fixed t , $Y^{(r)}(t)$ has a limit distribution as $r \rightarrow \infty$, after centering and norming. This relies on an elementary fact: if $\{N_n\}$ is a family of Poisson random variables with $E(N_n) \rightarrow \infty$ then

$$\frac{N_n - E(N_n)}{\sqrt{\text{Var}(N_n)}} \Rightarrow N(0, 1), \quad (n \rightarrow \infty). \quad (4.17)$$

From Eq. 4.3, we have

$$\begin{aligned} &P\left[\frac{Y^{(r)}(t) - b(r/t)}{a(r/t)} \leq x\right] = P(N([0, t] \times (a(r/t)x + b(r/t), \infty)) < r) \\ &= P\left[\frac{N([0, t] \times (a(r/t)x + b(r/t), \infty)) - tQ(a(r/t)x + b(r/t))}{\sqrt{r}}\right. \\ &\quad \left.< \frac{r - tQ(a(r/t)x + b(r/t))}{\sqrt{r}}\right). \end{aligned}$$

From Eq. 4.12, \sqrt{r} is asymptotic to the standard deviation of the Poisson random variable and so the left side random variable converges to a $N(0, 1)$ random variable.

Using (4.13), the right side converges to $\sqrt{t}h(x)$. We therefore conclude that under the left tail condition (4.11), for any fixed $t > 0$,

$$\lim_{r \rightarrow \infty} P \left(\frac{Y^{(r)}(t) - b(r/t)}{a(r/t)} \leq x \right) = \Phi(\sqrt{t}h(x)), \quad x \in \text{supp}_\gamma, \quad (4.18)$$

where $\Phi(x)$ is the standard normal cdf.

Now we can prove the following finite dimensional convergence.

Proposition 4.2 Assume (4.11) holds with $h(x)$ given in Eq. 4.16. Let $\{B(t), t \geq 0\}$ be standard Brownian motion. Then as $r \rightarrow \infty$,

$$\frac{Y^{(r)}(t) - b(r/t)}{a(r/t)} \Rightarrow h \leftarrow \frac{B(t)}{t}, \quad (4.19)$$

in the sense of convergence of finite dimensional distributions for $t > 0$.

Proof We illustrate the proof by showing bivariate pairs converge for two values of t . So suppose $0 < t_1 < t_2$ and $x_1 < x_2$ are in supp_γ and we show as $r \rightarrow \infty$,

$$\begin{aligned} P \left(\frac{Y^{(r)}(t_i) - b(r/t_i)}{a(r/t_i)} \leq x_i; i = 1, 2 \right) &\rightarrow P \left(h \leftarrow \frac{B(t_i)}{t_i} \leq x_i; i = 1, 2 \right) \\ &= P(B(t_i) \leq t_i h(x_i); i = 1, 2). \end{aligned} \quad (4.20)$$

We express the statements about $Y^{(r)}$ in terms of the Poisson counting measure and consider:

$$\begin{aligned} &\left(\begin{matrix} N([0, t_1] \times (a(r/t_1)x_1 + b(r/t_1), \infty)) \\ N([0, t_2] \times (a(r/t_2)x_2 + b(r/t_2), \infty)) \end{matrix} \right) \\ &= \left(\begin{matrix} N([0, t_1] \times (a(r/t_1)(x_1, x_2] + b(r/t_1), \infty)) + N([0, t_1] \times (a(r/t_1)x_2 + b(r/t_1), \infty)) \\ N([0, t_1] \times (a(r/t_2)x_2 + b(r/t_2), \infty)) + N([t_1, t_2] \times (a(r/t_2)x_2 + b(r/t_2), \infty)) \end{matrix} \right) \\ &= \left(\begin{matrix} N_1 + N_2 \\ N_3 + N_4 \end{matrix} \right) \end{aligned}$$

Consider the four terms N_i , $i = 1, \dots, 4$, in turn.

- (1) The term N_1 appropriately normed converges to 0:

$$\frac{N([0, t_1] \times (a(r/t_1)(x_1, x_2] + b(r/t_1), \infty)) - t_1 \Pi(a(r/t_1)(x_1, x_2] + b(r/t_1))}{\sqrt{r}} \Rightarrow 0. \quad (4.21)$$

The reason is that the centering is

$$\begin{aligned} \frac{t_1 \Pi(a(r/t_1)(x_1, x_2])}{\sqrt{r}} &= \frac{t_1 Q(ax_1 + b) - t_1 Q(ax_2 + b)}{\sqrt{r}} \\ &= \frac{r - t_1 Q(ax_2 + b)}{\sqrt{r}} - \frac{r - t_1 Q(ax_1 + b)}{\sqrt{r}} \\ &\rightarrow \sqrt{t_1}(h(x_2) - h(x_1)) > 0. \end{aligned}$$

So the left side of Eq. 4.21 is of the form $(N_r - \lambda_r)/\sqrt{r}$ where $\lambda_r/\sqrt{r} \rightarrow c > 0$ and thus

$$\text{Var } (N_r - \lambda_r)/\sqrt{r} = \lambda_r/r \rightarrow 0,$$

which verifies the convergence to 0 in Eq. 4.21.

- (2) The term N_2 becomes asymptotically normal. Let Z_1 be a standard normal random variable and apply (4.17) and (4.12) to get

$$\frac{N([0, t_1] \times (a(r/t_1)x_2 + b(r/t_1), \infty) - t_1 Q(a(r/t_1)x_2 + b(r/t_1)))}{\sqrt{r}} \Rightarrow \sqrt{t_1} Z_1.$$

- (3) For N_3 , despite its dependence on the variable t_2 , we also find

$$\frac{N([0, t_1] \times (a(r/t_2)x_2 + b(r/t_2), \infty) - t_1 Q(a(r/t_2)x_2 + b(r/t_2)))}{\sqrt{r}} \Rightarrow \sqrt{t_1} Z_1.$$

This result uses a combination of the reasoning that was used for N_1, N_2 .

- (4) The term N_4 is independent of N_1, N_2, N_3 so there is a standard normal variable $Z_2 \perp\!\!\!\perp Z_1$ and

$$\frac{N((t_1, t_2] \times (a(r/t_2)x_2 + b(r/t_2), \infty) - (t_2 - t_1) Q(a(r/t_2)x_2 + b(r/t_2)))}{\sqrt{r}} \Rightarrow \sqrt{t_2 - t_1} Z_2.$$

We conclude from this carving that

$$\left(\frac{N([0, t_1] \times (a(r/t_1)x_1 + b(r/t_1), \infty) - t_1 Q(a(r/t_1)x_1 + b(r/t_1)))}{\sqrt{r}} \right) \Rightarrow \left(\frac{\sqrt{t_1} Z_1}{\sqrt{t_1} Z_1 + \sqrt{t_2 - t_1} Z_2} \right),$$

as $r \rightarrow \infty$. Use (4.3) to write,

$$\begin{aligned} P\left(\left(\frac{Y^{(r)}(t_1) - a(r/t_1)}{b(r/t_1)}\right) \leq \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}\right) \\ = P\left(\left(\frac{N([0, t_1] \times (a(r/t_1)x_1 + b(r/t_1), \infty) - t_1 Q(a(r/t_1)x_1 + b(r/t_1)))}{\sqrt{r}}\right) \leq \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}\right) \end{aligned}$$

$$\begin{aligned}
&\leq \left[\frac{\left[\frac{-t_1 Q(a(r/t_1)x_1 + b(r/t_1))}{\sqrt{r}} \right]}{\left[\frac{-t_2 Q(a(r/t_2)x_2 + b(r/t_2))}{\sqrt{r}} \right]} \right] \\
&\rightarrow P(\sqrt{t_1}Z_1 \leq t_1 h(x_1), \sqrt{t_1}Z_1 + \sqrt{t_2 - t_1}Z_2 \leq t_2 h(x_2) \quad (\text{as } r \rightarrow \infty) \\
&= P\left(\frac{B(t_1)}{t_1} \leq h(x_1), \frac{B(t_2)}{t_2} \leq h(x_2)\right) \\
&= P \quad h^{\leftarrow}\left(\frac{B(t_1)}{t_1} \leq x_1, h^{\leftarrow}\left(\frac{B(t_2)}{t_2} \leq x_2 \right. \right.
\end{aligned}$$

This verifies (4.20). □

5 Final thoughts

The results of this paper suggest some obvious questions the answers to which have so far eluded us. Is there a jump process limit – presumably some sort of extremal process – in Eq. 4.19 corresponding to some sort of Poisson limit regime as opposed to the Brownian motion limit regime? In Proposition 4.2 is a stronger form of convergence – say in the J_1 -topology – possible? And so far, the mathematics of proving in a nice way that $\{Y^{(r)}, r \geq 1\}$ is Markov in the càdlàg space $D(0, \infty)$ has not cooperated.

References

- Arnold, B.C., Becker, A., Gather, U., Zahedi, H.: On the Markov property of order statistics. *J. Statist. Plann. Inference* **9**(2), 147–154 (1984)
- Cramer, E., Tran, T.H.: Generalized order statistics from arbitrary distributions and the Markov chain property. *J. Statist. Plann. Inference* **139**(12), 4064–4071 (2009)
- de Haan, L., Ferreira, A.: *Extreme Value Theory: An Introduction*. Springer, New York (2006)
- Deheuvels, P.: A Construction of Extremal Processes. In: *Probability and Statistical Inference* (Bad Tatzmannsdorf, 1981), pp. 53–57. Reidel, Dordrecht (1982)
- Deheuvels, P.: The strong approximation of extremal processes. II. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **62**(1), 7–15 (1983)
- Dwass, M.: Extremal processes. *Ann. Math. Statist* **35**, 1718–1725 (1964)
- Dwass, M.: Extremal processes. II. *Illinois. J. Math.* **10**, 381–391 (1966)
- Dwass, M.: Extremal processes. III. *Bull. Inst. Math. Acad. Sinica* **2**, 255–265 (1974). Collection of articles in celebration of the sixtieth birthday of Ky Fan
- Engelen, R., Tommassen, P., Vervaat, W.: Ignatov’s theorem: a new and short proof. *J. Appl. Probab.* **25A**, 229–236 (1988). A celebration of applied probability
- Goldie, C.M., Rogers, L.C.G.: The k -record processes are i.i.d. *Z. Wahrsch. Verw. Gebiete* **67**(2), 197–211 (1984)
- Goldie, C.M., Maller, R.A.: Generalized densities of order statistics *Statist. Neerlandica* **53**(2), 222–246 (1999)
- Ignatov, Z.: Ein von der Variationsreihe erzeugter Poissonscher Punktprozeß. *Annuaire Univ. Sofia Fac. Math. Méc.* **71**(2), 79–94 (1986). 1976/77
- Matheron, G.: *Random Sets and Integral Geometry*. Wiley, New York (1975). With a foreword by G.S. Watson, Wiley Series in Probability and Mathematical Statistics

- Molchanov, I.: Theory of Random Sets. Probability and its Applications (New York). Springer, London (2005)
- Rényi, A.: Théorie des éléments saillants d'une suite d'observations. Ann. Fac. Sci. Univ. Clermont-Ferrand No. **8**, 7–13 (1962)
- Resnick, S.I.: Tail equivalence and its applications. J. Appl. Probab. **8**, 136–156 (1971)
- Resnick, S.I.: Limit laws for record values. Stochastic Process. Appl. **1**, 67–82 (1973)
- Resnick, S.I.: Inverses of extremal processes. Adv. Appl. Probab. **6**, 392–406 (1974)
- Resnick, S.I.: Weak convergence to extremal processes. Ann. Probab. **3**(6), 951–960 (1975)
- Resnick, S.I.: Heavy Tail Phenomena: Probabilistic and Statistical Modeling. Springer Series in Operations Research and Financial Engineering. Springer, New York (2007). ISBN: 0-387-24272-4
- Resnick, S.I.: Extreme Values, Regular Variation and Point Processes. Springer, New York (2008). Reprint of the 1987 original
- Resnick, S.I., Rubinovitch, M.: The structure of extremal processes. Adv. Appl. Probab. **5**, 287–307 (1973)
- Rüschendorf, L.: Two remarks on order statistics. J. Statist. Plann. Inference **11**(1), 71–74 (1985)
- Shorrock, R.W.: On discrete time extremal processes. Adv. Appl. Probab. **6**, 580–592 (1974)
- Shorrock, R.W.: Extremal processes and random measures. J. Appl. Probab. **12**, 316–323 (1975)
- Stam, A.J.: Independent Poisson processes generated by record values and inter-record times. Stochastic Process. Appl. **19**(2), 315–325 (1985)
- Vervaat, W., Holwerda, H. (eds.): Probability and Lattices, Volume 110 of CWI Tract. Stichting Mathematisch Centrum, Centrum voor Wiskunde en Informatica, Amsterdam (1997)
- Weissman, I.: Extremal processes generated by independent nonidentically distributed random variables. Ann. Probab. **3**, 172–177 (1975)